

Study of Newly Synthesized Pyridinium-based Cationic Surfactants for Drug Interaction and Antibacterial Activity

Ali Jaan^{1,2}, Saqib Ali², Mohsin Javed¹, Ali Haider², Khurram Shahzad Munawar^{3,4*},
Saja Abdulrahman Althobaiti⁵, and Mahboob ur Rehman⁶

¹Department of Chemistry, School of Science, University of Management and Technology,
C-II Johar Town, Lahore, Pakistan

²Department of Chemistry Quaid-i-Azam University, 45320, Islamabad, Pakistan

³Institute of Chemistry, University of Sargodha, Sargodha, 40100, Pakistan

⁴Department of Chemistry, University of Mianwali, Mianwali, 42200, Pakistan

⁵Department of Chemistry College of Sciences and Humanities, Prince Sattam Bin
Abdulaziz University, Saudi Arabia

⁶Cardiology Department, Pakistan Institute of Medical Sciences (PIMS), Islamabad, Pakistan




Fig. S1. FT-IR spectrum of sample A1.

* Corresponding Author: Khurram Shahzad Munawar <khurramchemist@gmail.com>





Fig. S2. FT-IR spectrum of sample A2.

Fig. S3. ¹H-NMR spectrum of sample A1.




Fig. S4. ¹³C-NMR spectrum of sample A1.




Fig. S5. ¹H-NMR spectrum of sample A2.




Fig. S6. ^{13}C -NMR spectrum of sample A2.




Fig. S7. Structure of ketoprofen.




Fig. S8. Inhibition zones of samples A1 and A2 against *Staphylococcus pseudo*.




Fig. S9. Inhibition zones of samples A1 and A2 against *Klebsiella pneumoniae*.




Fig. S10. Inhibition zones of samples A1 and A2 against *Staphylococcus aureus*.